



Evaluation of Pakistani Wheat Cultivars for Yield-Related Traits and Yellow Rust Resistance in Agro-Climatic Conditions of Dera Ismail Khan

Inayat Ullah¹(Corresponding Author), Nimra Qayyum², Zain Ul Abidin³, Muhammad Yousaf Shah Hashmi⁴, Fayaz Ullah⁶, Muhammad Raza⁷

¹ MSc Hons student, Department of Plant Breeding & Genetics, Gomal University Dera Ismail Khan, Pakistan. Email: inayatkhanbehran445@gmail.com

^{2,3} MSc Hons student, Department of Plant Breeding & Genetics, Gomal University Dera Ismail Khan, Pakistan

^{4,6,7} MSc Hons student, Department of Plant Breeding & Genetics, Gomal University Dera Ismail Khan, Pakistan

Abstract

The primary crop for feeding the planet's expanding population is wheat (*Triticum aestivum* L.), which is still a common ingredient in many cuisines. The breeding and development of semi-dwarf and high-yielding varieties have boosted national wheat yield. The constantly changing climate necessitates to development of more climate-resilient wheat varieties that can better withstand disease pressure, heat and drought stress. This study was designed to evaluate elite Pakistani wheat cultivars for yield-related traits and yellow rust resistance in the agro-climatic conditions of Dera Ismail Khan. Twenty-six Pakistani wheat varieties were grown in a standardized design as a Randomized Complete Block Design (RCBD) using three replications. Data collected for days to 50% heading, plant height, spike length, 1000GW and yellow rust resistance was subjected to the R package for statistical analysis. AZRC Dera was found to have the earliest heading (104 days). Gulzar-19 was found to be the most dwarf variety (mean height 68cm) while Peer Sabaq-19 was the tallest one (mean height 100 cm). Mean spike length ranged from 8.8 cm to 15 cm. The highest spike length was noted for NIFA Lalma-13 (15 cm) and the lowest for Faisalabad-2008 (8.8 cm). The highest 1000GW was recorded for the recently developed variety of ARI-RK2022 (65g), while the lowest 1000GW was recorded for rust susceptible wheat variety Faisalabad Punjab (40g). Based on the yellow rust score scale, yellow rust highly resistant (HR) cultivars were AZRC Dera, Khaista, Wadan, Akbar19, Shahkar, and Gulzar19. Moderately resistant cultivars (MR) include Hashim08, Shahid17, Fakhr-e-Bahakkar, and Tarnab Gandam. Moderately susceptible (MS) cultivars include Shalimar-88, while highly susceptible (HS) cultivars were Faisalabad Punjab, Faisalabad-2008, and Punjab-1. Based on the findings of this study, AZRC Dera, RK2022 and NIFA Lalma-13 are recommended to be grown by the local farmers in the agro-climatic conditions of Dera Ismail Khan.

Keywords: Wheat yield, yellow rust, spike length, Days to 50% heading, 1000GW

DOI: <https://zenodo.org/records/14913568>

Journal Link: <https://jai.bwo-researches.com/index.php/jwr/index>

Paper Link: <https://jai.bwo-researches.com/index.php/jwr/article/view/101>

Publication Process Received: 24 Jan 2024/ Revised: 15 August 2024/ Accepted: 20 Sep 2024/ Published: 23 Sep 2024

ISSN: Online [3007-0929], Print [3007-0910]

Copyright: © 2025 by the first author. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Indexing:



Publisher: BWO Research International (15162394 Canada Inc.) <https://www.bwo-researches.com>

Introduction

The primary crop for feeding the planet's expanding population is wheat (*Triticum aestivum* L.), which is still a common ingredient in many cuisines. Around the world, it is grown on more than 220 million hectares, with an annual yield of more than 749 million tonnes. The tetraploid domesticated wheat *T. turgidum* ssp. *dicoccum*, which contributed the AA and BB sub-genomes, and the wild grass species *Aegilops tauschii*, which contributed the DD sub-genome, naturally hybridized to produce the hexaploid bread wheat ($2n = 6x = 42$, genome AABBDD (Peterson et al., 2006). Allohexaploid bread wheat (*Triticum aestivum* L.) is a species that resulted from two subsequent rounds of hybridization. It is believed that the second hybridization event took place between 8000 and 10,000 years ago in the Fertile Crescent (Heun et al., 1997; Salamini et al., 2002). The size of the wheat genome is around 17 Gbps, and it is very complex, especially in terms of chromosomal duplications and rearrangements and the extremely high proportion of repetitive sequences (Akpınar et al., 2015).

The germplasm of bread wheat has thus changed throughout historic human migratory routes. From 8500 to 2300 years ago, the earliest farmers from this region carried it both westward to Europe and eastward to Asia, and domesticated wheat populations have spread over Europe and Asia, where they have adapted to local conditions to become known as landraces. The introduction of bread wheat to the New World began in the 16th century, first in Latin America, then in North America and Australia (Stevens et al., 2016). According to the experts, they were able to follow the evolution of wheat from the Fertile Crescent to the regions of Europe and Asia where it was planted and raised. They point out that the Green Revolution

caused a significant alteration in wheat that diminished its variety. According to their research, the majority of cultivars planted now are descended from strains that were created in southeastern Europe around the Mediterranean Sea and on the Iberian Peninsula. They point out that the lack of diversity creates a chance for further advancements in wheat production. Asian wheat varieties, they point out, are a good source of variation and may be utilized in the next research projects targeted at boosting crop yields as a result (Balfourier et al., 2019).

Wheat is the main staple food item of the country's population and the largest grain crop as well. Pakistan has been regularly importing wheat, except for a few years. Wheat's domestic production has remained short of its domestic demands. To estimate the wheat varietal effect on wheat production. The development of modern wheat varieties represents an advancement in agricultural science driven by the need for food security and crop resilience (Shiferaw et al., 2013; Abraham et al., 2014; Mondal et al., 2016). This process integrates traditional breeding techniques with cutting-edge genetic engineering to produce wheat that is more resistant to disease, pests, and environmental stresses. Modern varieties are designed to achieve higher yields, improved nutritional quality, and greater adaptability to changing climatic conditions. Through meticulous selection and hybridization, researchers can introduce desirable traits from diverse wheat lines, accelerating the creation of superior cultivars. This ongoing innovation not only boosts productivity but also ensures sustainable farming practices, contributing to the global effort to meet the dietary needs of a growing population (Razzaq et al., 2021; Singha and Sangha, 2024). The current study was designed to

evaluate elite Pakistani wheat cultivars for yield-related traits and yellow rust resistance in the agro-climatic conditions of Dera Ismail Khan.

Materials & Methods

The current research was conducted in an experimental area of Gomal University, Dera Ismail Khan, in the wheat growing season from November 2023 to May 2024.

Research Design

The research design for this study was standardized as RCBD (randomized complete block design) using three replications in the design. The experimental plot area was kept at 18 m × 5 m (90 m²) with 26 wheat germplasm replication⁻¹ having a size of, each line being kept 5 meters in length and a 30 cm distance between lines. Sowing was carried out in lines, and ten seeds per row were dibbled, keeping a plant-to-plant distance of 15 cm and row-to-row distance standardized at 30 cm.

Practices

The land was prepared using 2-3 deep ploughings accompanied by rotavator operation. There was application of regular necessary agricultural practices such as fertilizer application of recommended doses of nitrogen and phosphorous, irrigation, hoeing etc.

Material

The germplasm comprising 26 wheat varieties was grown in the wheat growing season of 2022-2023 (Table 1). All 26 wheat varieties were acquired from the National Agriculture Research Council (NARC), Islamabad.

Table 1: Germplasm used in this study.

S/No	Germplasm name	S/No	Germplasm name
1	Hashim-08	14	PS-15
2	AZRC Dera	15	PS-19
3	Khaista	16	Punjab-11
4	Wadan-17	17	Shalimar-88
5	Akbar-19	18	Zamindar-80

6	Akbar Punjab	19	Peersabaq-15
7	Shahkar	20	Peersabaq-19
8	Gulzar-19	21	Peersabaq-21
9	Shahid-17	22	Khyber-23
10	Peer Sabaq-13	23	Tarnab Gandam
11	Fakhre Bakkar	24	Taskeen-22
12	Faisalabad Punjab	25	NIFA Lalma-13
13	Faisalabad 2008	26	RK-2022

Data collection

Data was recorded keeping selection intensity at 5% from every line. The protocol for data collection of parameters is as under.

Parameters

1. Days to 50% heading

Days to 50% heading was documented from the time of sowing of the crop to the occasion of 50% heading in a specific line.

2. Yellow rust

The scoring method of leaf rust considered the percentage of leaf area covered by pustules on the most heavily infested leaves, using a scale from 0 to 100.

3. Plant height

Plant height was calculated by measuring the distance from the base of the main culm to the tip of the spike awn of the selected tiller.

4. Spike length (cm)

The spike length of the randomly chosen plants was calculated after measuring the spike from the base to the pinnacle of the spike, exclusive of awns and further analyzed for their average.

5. 1000-grain weight (g)

At the stage of physiological maturity, 1000 grains were calculated from the randomly selected plants and their weight was taken in grams.

Statistical analysis

The data collected for yield-related traits was subjected to the R package for graphical presentation.

Results & Discussion

Days to 50% heading

Days to 50% heading is an important yield-related attribute which determines the early seed setting, early maturity and greater yield potential. Wheat cultivars requiring maximum days to 50% heading face extreme terminal heat stress during anthesis and maturity, thereby experiencing late maturity and reduced yield, especially when late-sown or in heat stress-prone areas like Dera Ismail Khan. Modern breeding cultivars are developed for early maturity to bypass heat stress during the early stages of anthesis and maturity. Our results showed that all cultivars showed differences for days to 50% heading. AZRC Dera was found to have the earliest heading (104 days) while Wadan-17 and Peer Sabaq-19 were found to be the late heading ones (116 days) as shown in Fig. 1. Our findings are supported by [Nahar et al., \(2010\)](#) who showed that in case of late seeding, the varieties phased a significant level of high-temperature stress that also significantly affected the required days to germination, booting, anthesis, maturity of all varieties including the yield as compared to normal sowing treatment. The temperature during the grain filling or grain maturing period in their trial was near 23°C in the case of normal seeding, and it was near 28°C to 30°C and sometimes reached above this range in the later period of late-seeded treatment. In the normal sowing treatment, the germination period was lower than the late sowing treatment, as during that time the temperature was higher as compared to the late sowing condition, where the temperature was lower. Days to anthesis and booting decreased due to late sown heat stress conditions, regardless of the cultivars. These phenological characteristics under heat-stressed conditions led the wheat cultivars to

significantly lower the grain yield as compared to normal conditions. Due to heat stress, the yield reduction was 69.53% in the wheat variety 'Sourav', 58.41% in 'Pradip', 73.01% in 'Sufi', 55.46% in 'Shatabdi' and 53.42% in 'Bijoy'.




Fig. 1 Variations among germplasm for days to 50% heading. Black arrows show early and late-heading germplasm.

Resistance to yellow rust

Yellow/stripe rust of wheat, caused by *Puccinia striiformis* f. sp. *tritici* (Pst), is an important wheat foliar disease, mostly common in hot, humid wheat growing regions and causes heavy yield losses during epidemics. Severe infestation caused by yellow rust has been reported both in Pakistan ([Ali et al., 2009](#)) and elsewhere ([Singh et al., 2004](#)). In Pakistan, almost 70% of the acreage under wheat is prone to the disease. This biotroph reduces the total photosynthetic area, utilizes the plant's assimilates and interrupts with normal growth of the host, leading to a reduction in yield. Tremendous variations for yellow rust resistance were observed among all germplasm tested in this study. Yellow rust highly resistant (HR) cultivars include AZRC Dera, Khaista, Wadan, Akbar19, Shahkar, and Gulzar19. Moderately resistant cultivars (MR) include Hashim08, Shahid17, Fakhr-e-Bahakkar, and Tarnab Gandam. Moderately susceptible (MS) cultivars

include Shalimar-88, while highly susceptible (HS) cultivars include Faisalabad Punjab, Faisalabad-2008, and Punjab-1, as shown in Fig. 2.




Fig. 2 Variations among germplasm for yellow rust resistance.

3. Plant height

No obvious differences were observed in height among cultivars, the reason being the exploitation of semi-dwarf nurseries of CIMMYT in wheat breeding programs in Pakistan. Gulzar-19 was a very dwarf variety (mean height: 68cm) while Peer Sabaq-19 was the tallest one (mean height: 100 cm) as shown in Fig. 3. Plant height is a significant trait influencing wheat yield due to its impact on lodging resistance and biomass accumulation. The height of wheat plants can vary widely among different varieties cultivated in Pakistan. Plant height is an important trait affecting wheat yield as it influences lodging resistance and biomass accumulation. Studies have shown significant variation in plant height among different wheat varieties. Taller varieties like 'Inqalab-91' and 'Punjab-96' have historically been popular due to their higher biomass production, which is useful for fodder. Varieties such as 'Seher-06' and 'NARC-2009' are known for their short stature, which enhances lodging resistance and is beneficial in regions prone to heavy winds and rain (Mumtaz et al., 2015).




Fig. 3 Variations among germplasm for plant height.

4. Spike length

The grain yield of wheat is a variable trait that depends mainly on yield components and environmental factors. However, the variability of yield is more studied than yield components. The investigation of variability and components of phenotypic variance for plant height and spike length is very important for the cultivar creation in breeding programs. These two traits are quantitative characteristics which are in correlation with other yield components. Significant variations were found for spike length among all wheat varieties. Mean spike length ranged from 8.8 cm to 15 cm. The highest spike length was noted for NIFA Lalma-13 (15 cm) and the lowest for Faisalabad-2008 (8.8 cm). Our findings are supported by Zečević et al. (2008). They investigated genetic and phenotypic variability for plant height and spike length in ten winter wheat cultivars from different selection centres. The experiment was performed in a randomized block design in five replications on the experimental field in two years. A total of 50 plants have been analyzed in the full maturity stage. Average estimated values of plant height and spike length differed significantly among years and cultivars. The highest average value for plant height was established at the Ljubičevka cultivar

(89cm), and the lowest average plant height was established at Zagrepčanka (64.0cm). Higher values were established in the second year than in the first year of investigation (69.1cm and 63.9cm, respectively). The variation coefficient of plant height for all examined cultivars and years varied from 2.7% to 6.1%. The highest average value for spike length was Ljubičevka (9.8cm), and the lowest was Slavonija (7.0cm).




Fig. 4 Variations among germplasm for spike length.

1000GW

1000-grain weight (1000GW) is one of the most important determinants of wheat grain yield and is determined by grain length (GL), grain width (GW) and grain thickness. This quantitative trait is highly affected by environmental factors, especially when wheat germplasm is grown later than the normal time, which later experiences heat stress at the grain filling stage, thereby reducing the 1000-grain weight. In our study, the highest 1000GW was recorded for the recently developed variety of ARI- RK2022 (65g), followed by NIFA Lalma-13 (60g) while the lowest 1000GW was recorded for rust susceptible wheat variety Faisalabad Punjab (40g) as shown in Fig. 5. Our results are by [Dhyani et al., \(2013\)](#) who performed who evaluated wheat genotypes viz., DBW-140, Raj-3765, PBW-574, K-0-307 and

HS-240 under timely/late sown and heat stress conditions for 1000GW and other yield-related traits. Their results showed that the heat susceptibility index (HIS) for 1,000-grain weight, grain weight and grain yield of wheat genotypes viz., HS 240 and K-0-307 was highest as compared with DBW 140, Raj 3765 and PBW 574 genotypes.




Fig. 5 Variations among germplasm for 1000GW. Black arrows show germplasm with the lowest and highest 1000GW.

Conclusion & Recommendations

All 26 cultivars used in this study showed significant differences for different yield-related traits except plant height. AZRC Dera was found to have the earliest heading (104 days). Gulzar-19 was found to be the most dwarf variety (mean height 68cm) while Peer Sabaq-19 was the tallest one (mean height 100 cm). Mean spike length ranged from 8.8 cm to 15 cm. The highest spike length was noted for NIFA Lalma-13 (15 cm) and the lowest for Faisalabad-2008 (8.8 cm). The highest 1000GW was recorded for the recently developed variety of ARI-RK2022 (65g), while the lowest 1000GW was recorded for rust susceptible wheat variety Faisalabad Punjab (40g). Based on the yellow rust score scale, yellow rust highly resistant (HR) cultivars were AZRC Dera, Khaista, Wadan, Akbar19, Shahkar, and Gulzar19. Moderately resistant cultivars (MR)

include Hashim08, Shahid17, Fakhr-e-Bahakkar, and Tarnab Gandam. Moderately susceptible (MS) cultivars include Shalimar-88, while highly susceptible (HS) cultivars were Faisalabad Punjab, Faisalabad-2008, and Punjab-1. Based on the findings of this study, AZRC Dera, RK2022 and NIFA Lalma-13 are recommended to be grown by the local farmers in the agro-climatic conditions of Dera Ismail Khan.

Authors Contribution

Nasr Ullah Khan conceived the idea, designed the study and drafted the manuscript. Inayat Ullah, Nimra Qayyum and Zain Ul Abidin conducted the experiments and collected the data. Muhammad Yousaf Shah Hashmi, Fayaz Ullah and Muhammad Raza helped in data collection and data analysis.

References

- Abraham B, Araya H, Berhe T, Edwards S, Gujja B, Khadka RB, Koma YS, Sen D, Sharif A, Styger E, Uphoff N. The system of crop intensification: reports from the field on improving agricultural production, food security, and resilience to climate change for multiple crops. *Agriculture & Food Security*. 2014; 3:1-2
- Akpinar BA, Kantar M, Budak H. Root precursors of microRNAs in wild emmer and modern wheat show major differences in response to drought stress. *Functional & integrative genomics*. 2015; 15:587-98
- Ali S, Shah SJ, Rahman HU, Saqib MS, Ibrahim M, Sajjad M. Variability in wheat yield under yellow rust pressure in Pakistan. *Turkish Journal of Agriculture and Forestry*. 2009; 33(6):537-46
- Balfourier F, Bouchet S, Robert S, De Oliveira R, Rimbart H, Kitt J, Choulet F, International Wheat Genome Sequencing Consortium, BreedWheat Consortium, Paux E. Worldwide phylogeography and history of wheat genetic diversity. *Science advances*. 2019; 5(5):eaav0536
- Dhyani K, Ansari MW, Rao YR, Verma RS, Shukla A, Tuteja N. Comparative physiological response of wheat genotypes under terminal heat stress. *Plant signalling & behaviour*. 2013 Jun 1;8(6):e24564.
- Heun M, Schafer-Pregl R, Klawan D, Castagna R, Accerbi M, Borghi B, Salamini F. Site of einkorn wheat domestication identified by DNA fingerprinting. *Science*. 1997; 278(5341):1312-4
- Mondal S, Rutkoski JE, Velu G, Singh PK, Crespo-Herrera LA, Guzman C, Bhavani S, Lan C, He X, Singh RP. Harnessing diversity in wheat to enhance grain yield, climate resilience, disease and insect pest resistance and nutrition through conventional and modern breeding approaches. *Frontiers in Plant Science*. 2016; 7:991
- Mumtaz MZ, Aslam M, Nasrullah HM, Akhtar M, Ali B. Effect of various sowing dates on growth, yield and yield components of different wheat genotypes. *American-Eurasian Journal of Agricultural and Environmental Science*. 2015;15(11):2230-4
- Nahar K, Ahamed KU, Fujita M. Phenological variation and its relation with yield in several wheat (*Triticum aestivum* L.) cultivars under normal and late sowing mediated heat stress conditions. *Notulae Scientia Biologicae*. 2010; 2(3):51-6
- Peterson JJ, Beecher GR, Bhagwat SA, Dwyer JT, Gebhardt SE, Haytowitz DB, Holden JM. Flavanones in grapefruit, lemons, and limes: A compilation and review of the data from the analytical literature. *Journal of food composition and analysis*. 2006; 19:S74-80
- Razzaq A, Kaur P, Akhter N, Wani SH, Saleem F. Next-generation breeding strategies for climate-ready crops. *Frontiers in Plant Science*. 2021; 12:620420
- Salamini F, Özkan H, Brandolini A, Schäfer-Pregl R, Martin W. Genetics and geography of wild cereal domestication in the Near East. *Nature Reviews Genetics*. 2002; 3(6):429-41
- Shiferaw B, Smale M, Braun HJ, Duveiller E, Reynolds M, Mauricio G. Crops that feed the world 10. Past successes and future challenges to the role played by wheat in global food security. *Food security*. 2013; 5:291-317
- Singh RP, William HM, Huerta-Espino J, Rosewarne G. Wheat rust in Asia: meeting the challenges with old and new technologies. In *Proceedings of the 4th International Crop Science Congress 2004*; 26: 1-13.
- Singha S, Singha R. Crop Improvement Strategies and Principles of Selective Breeding. In *Water-Soil-Plant-Animal Nexus in the Era of Climate Change*. 2024; 93-113. IGI Global
- Stevens CJ, Murphy C, Roberts R, Lucas L, Silva F, Fuller DQ. Between China and South Asia: A Middle Asian corridor of crop dispersal and agricultural innovation in the Bronze Age. *The Holocene*. 2016; 26(10):1541-55
- Zečević V, Knežević D, Mićanović D, Madić M. Genetic and phenotypic variability of spike length and plant height in wheat. *Kragujevac Journal of Science*. 2008; 30:125-30.