



International Journal of Agriculture Innovations and Cutting-Edge Research



Response of Different Concentrations of Liquid Pollen Applications on Fruit Set, Yield and Fruit Physical Properties of Dhakki and Zahidi Date Palm (*Phoenix dactylifera* L.) under the Agro-Ecological Conditions of Dera Ismail Khan

Bilal Ahmad Khan¹, Muhammad Jawad Nazir²(Corresponding Author), Rashid Khan³, Niaz Ud Din⁴, Ehtesham Ul Haq⁵

- ¹ PhD scholar, Faculty of Agriculture, Department of Horticulture, Gomal University, DIKhan, Pakistan, ORCID: <https://orcid.org/0009-0006-4273-520X>, Email: Bilal.hort@gmail.com
- ² Researcher, Plant Genetic Resources Institute (PGRI), National Agri Research Centre, Islamabad, Pakistan, ORCID: <https://orcid.org/0009-0001-9042-9614>, Email: jawadnazir1997@gmail.com
- ³ MPhil scholar, Department of Agronomy, Faculty of Agriculture, Gomal University, DIKhan, Pakistan, ORCID: <https://orcid.org/0009-0009-6213-3048>, Email: rrs125634@gmail.com
- ⁴ MPhil scholar, Faculty of Agriculture, Department of Horticulture, Gomal University, DIKhan, Pakistan, ORCID: <https://orcid.org/0009-0004-2843-2152>, Email: niazuddin4373@gmail.com
- ⁵ PhD scholar, Department of Agronomy, Faculty of Agriculture, Gomal University, DIKhan, Pakistan, ORCID: <https://orcid.org/0009-0001-2574-4722>, Email: malikehteshamdhap@gmail.com

Abstract

The main cause for low fruit set %, retaining ability, fruit weight and fruit yield per plant is the improper pollination in most of the commercial date palm varieties in different date palm growing countries of the world. Palms were fertilized by the suspended pollen spray method. Results of this study revealed that the maximum days to fruit set (23.40 and 22.10), fruit set (96.47 and 91.46%), fruit drop (43.7 and 26.81%), bunch weight (11.13 and 6.41 kg), and fruit yield (178.13 and 64.13 kg plant⁻¹) was recorded with the maximum pollen strength of 8 g L⁻¹ in both the cultivars Dhakki and Zahidi, respectively. On the other hand, the minimum values of these parameters were shown by the pollen strength of 2 g L⁻¹ in both the date palm cultivars. Overall, the cultivar Dhakki achieved maximum values of fruit quality traits, with fruit size, fruit weight, and pulp weight. Hence, based on this study, the liquid pollen application of 8 g L⁻¹ is suggested for attaining maximum fruit set and yields in date palm under the Agroecological situations of Dera Ismail Khan.

Keywords: Date palm, Varieties, Dhakki, Zahidi and pollination.

DOI: <https://zenodo.org/records/16792835>

Journal Link: <https://jai.bwo-researches.com/index.php/jwr/index>

Paper Link: <https://jai.bwo-researches.com/index.php/jwr/article/view/146>

Publication Process Received: 14 Jul 2025/ Revised: 08 Aug 2025/ Accepted: 10 Aug 2025/ Published: 12 Aug 2025

ISSN: Online [3007-0929], Print [3007-0910]

Copyright: © 2024 by the first author. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Indexing:



Publisher:

BWO Research International (15162394 Canada Inc.) <https://www.bwo-researches.com>

Introduction

The date palm (*Phoenix dactylifera* L.) belongs to the family Arecaceae and is distributed in tropical and sub-tropical countries of the world. It is commonly commercially cultured in the arid areas of the Middle East and North Africa (Henderson, 2009). Dates are eaten in dried or fresh form. They are also used for the preparation of salads, syrups, cakes, etc. Date fruit contains a high number of calcium, magnesium, phosphorus, potassium, iron, zinc, copper, manganese, selenium, vitamins A, A1, B, B1, B2, B3, B5, B6, and C, as well as a variety of amino acids (Anon, 2006). Thiamine, riboflavin, niacin, and pantothenic acid are also found in dates. These vitamins and minerals aid in the production of hemoglobin, a protein found in red blood cells that binds to oxygen and transports it from the lungs to the body's tissues (Anon, 2005). Dates also deliver 3 g of dietary fibre and 29 g of natural sugars with fructose, glucose, and sucrose. To put it another way, one date has 31 g of carbs, which delivers the body with a lot of energy (Anon, 2007).

The pollen from date palm (*Phoenix dactylifera*) blossoms has been demonstrated to contain estrone, a form of estrogen with notable gonadotropic effects observed in young rats. Recent experimental studies have confirmed that date palm pollen extract can elevate circulating levels of estradiol, luteinizing hormone (LH), and follicle-stimulating hormone (FSH) in both male and female rats, underscoring its potential endocrine modulating properties (Abdelrahim et al., 2015; Moshfegh et al., 2016).

Pakistan ranks in the 6th position in the top 10 countries producing with an annual production of 524,286 tonnes (FAO, 2017). In Pakistan, date palm is commercially grown in all tropical and subtropical areas of the country. Pakistan spreads fresh and dried

dates to other countries and earns billions of rupees annually (Maryam et al., 2015). After Tunisia, Pakistan is the second-largest exporter (9,781 tonnes). The major distributors of Pakistani dates are India, Bangladesh, Germany, Italy, Germany, Japan, Canada, Denmark, UAE, Australia, South Africa, Saudi Arabia, the United Kingdom, and the United States.

The utmost common technique of pollination is being used by date growers are 1) to cut the strands of male at maturity stage when pollen can fertilize female flowers and placed into the female spathe, 2) extracted dried pollen are suspended and sprayed on open spathe, 3) dusting of full fresh male spathes or extracted dried pollens are dusted by duster, 4) extracted dried pollens are rubbed into cotton bowl and placed into open spathes of female plant. Various past studies have elaborated on the impact of different pollination techniques on the fruit set and yield of date palm. Iqbal et al. (2010) investigated the effects of different pollination techniques, i.e., placement, dusting and spray of diluted pollens on Dhakki date. Harash and Abdul Naser (2010) sprayed liquid pollen on Khalas date clusters and found that the spray approach yielded the best yield when associated with other treatments. Abdalla et al. (2011) pollinated cv. Zaghloul used the conventional approach (placing male strands) as well as pollen spray and boron. The maximal fruit set and weight were measured using the (strand placement method). Abdullah et al. (2014) observed two pollination techniques for the cv. Khalas date palm: rubbing pollen on a sponge and inserting male threads inside the fruit, and found no important changes in fruit set, yield, or morphologic features of the fruit in either strategy. Samouni et al. (2016) pollinated Saidy date palms using dusting methods (mixture of pollen +

starching at different concentrations with active ingredient) and traditional methods (placement of 5 male strands/spathe) and found that dusting methods resulted in the highest fruit set, yield, and quality when compared to traditional methods. The impact of pollination with suspended pollen grains spray on the Segue cv of date palm was investigated by Soliman et al. (2017). Traditional pollination was used, which included placing 5 male strands/spathe, spraying 3 g pollen L⁻¹, 3 g pollen + 2 g sugar per L⁻¹, 2 g pollen L⁻¹, and 2 g pollen + 3 g sugar L⁻¹ are all options. In terms of yield and yield components, traditional pollination was shown to be superior. Soliman et al. (2017). Impact of pollination by pollen-grain water suspension spray on yield and fruit quality of the Segae date palm cultivar (*Phoenix dactylifera* L.). *Pakistan Journal of Botany*, 49(1), 119–123. This study reported that spray-pollination with a mix of 2 g pollen + 3 g sugar per litre significantly increased total sugar percentage, and 2 g pollen per litre led to higher acidity percentages in date fruits.

Problem Statement

Date palm (*Phoenix dactylifera* L.) is a culturally and economically vital fruit crop in arid and semi-arid regions, particularly in southern Khyber Pakhtunkhwa, Pakistan. Among the locally cultivated varieties, Dhakki and Zahidi are commercially important due to their high market value and consumer preference. However, suboptimal pollination, especially under fluctuating weather and manual labour constraints, remains a major bottleneck in achieving consistent fruit set and optimal yield. Conventional hand pollination practices are time-consuming labor labour-intensive, and often fail to ensure uniform pollen distribution, leading to poor fertilization and reduced fruit quality. In this context, liquid pollen application has emerged as a promising

alternative that can improve pollination efficiency and reduce dependency on traditional methods.

Despite its potential, limited scientific data exist on the impact of different concentrations of liquid pollen sprays on fruit set, yield attributes and physical fruit characteristics of leading cultivars like Dhakki and Zahidi. Moreover, no such work has been reported specifically under the agroecological conditions of Dera Ismail Khan, a region with unique climatic features that can influence pollen viability, stigma receptivity and fruit development. This research addresses this critical knowledge gap by systematically evaluating how varying concentrations of liquid pollen affect reproductive success and post-pollination fruit quality traits.

Objectives of the Study

1. To evaluate the effect of different concentrations of liquid pollen spray on fruit set percentage in Dhakki and Zahidi date palm cultivars.
2. To determine the influence of liquid Pollen concentrations on fruit yield per bunch and per plant under field conditions.
3. To assess the changes in fruit physical traits (length, diameter, weight, pulp-to-seed ratio) in response to liquid pollen applications.
4. To compare the varietal responses (Dhakki vs. Zahidi) to liquid pollen treatments under the climatic conditions of Dera Ismail Khan.
5. To propose optimized pollen management strategies for improving pollination efficiency and productivity in commercial date palm cultivation.

Materials and Methods

The trial took place in Gomal University's Date Palm Research Orchard in Dera Ismail Khan, Pakistan, from March to August 2020. Four trees of each Dhakki and Zahidi date palm, taking similar age

and vigorous growth, were selected. The trial was conducted utilizing split plot configurations in R.C.B.D. Female cultivars were retained as the primary factor, pollen concentrations as a sub-factor, and pollination was done with a pollen source three times. In the month of March, pollen was collected. Male stamens were collected at the maturity phase and carefully transported to the laboratory. The protective sheath was removed, and these inflorescences were placed on newspaper for the extraction of pollen under the sun. After 2-3 hours, the flowers were opened, and pollen grains were released. Pollens were extracted by shaking the spathes. These pollens were dried in the sun for 1-2 hours by spreading them on a newspaper. These pollens were dried and then meshed with a porous cloth to remove any undesirable elements before being packed in an airtight glass bottle and stored at 4 °C in a household refrigerator. The number of these pollen 2, 4, 6 and 8 (g) was suspended in one litre of refined water. Three spathes of each female were sprayed by hand spray pump at noon to avoid natural pollination, and bagging was done after pollination. Throughout the research, cultural actions like as hoeing, watering, and weeding were all carried out equally.

Showing the details of treatments

Pollination treatments	Varieties	
	Dhakki	Zahidi
T ₁	sprig of 2 g pollen/litre of water	sprig of 2 g pollen/litre of water
T ₂	sprig of 4 g pollen/litre of water	sprig of 4 g pollen/litre of water
T ₃	sprig of 6 g pollen/litre of water	sprig of 6 g pollen/litre of water
T ₄	sprig of 8 g pollen/litre of water	sprig of 8 g pollen/litre of water

Statistical Analysis

The experimental data collected on fruit set percentage, yield per bunch and per plant, as well as various physical fruit traits (fruit length, diameter, weight, pulp-to-seed ratio), were subjected to statistical analysis using SPSS version 26.0 and Microsoft Excel 2019.

The analysis of variance (ANOVA) was performed to determine the significance of treatment effects across different concentrations of liquid pollen for both Dhakki and Zahidi cultivars. Means were separated using the Least Significant Difference (LSD) test at 5% level of significance ($p \leq 0.05$) to compare varietal responses and pollen concentration impacts.

Results and Discussion

Fruit set (%)

Table 1 indicated that several pollen concentrations had a significant influence on fruit set percent, whereas date varieties and contact were determined to be non-significant. The pollen concentration showed that the highest fruit set (93.9%) was observed in T₄ (8 g L⁻¹), followed by T₃ (6 g L⁻¹) and T₂ (4 g L⁻¹), by making fruit set % (89.8 and 74.9%, respectively). Although the lowest fruit set (62.8%) was noted in T₁ (2 g L⁻¹). The highest fruit set in T₄ (8 g L⁻¹) might be due to the maximum number of pollens present in the solution, which fertilized the maximum number of female flowers (fruit set). These results are in agreement with the results of [Munir et al. \(2019\)](#), who noted the highest fruit set in cv. Zahidi and Khudrawi at a higher pollen concentration of 4 g L⁻¹ instead of 1 g L⁻¹. Similarly, [Zahara et al. \(2019\)](#) reported the highest fruit set in cultivar Barhai and Medjhol when pollinated with higher concentrations of suspended pollen.

Days to fruit set

The data in Table 1 showed that the pollen concentration caused significant

variation in days to fruit set, though date varieties and their interaction could not produce any significant result regarding days to fruit set. The maximum days to fruit set (22.75 days) were counted in T₄ (8 g L⁻¹), followed by T₃ (6 g L⁻¹) and T₂ (4 g L⁻¹), which gave 21.02 and 19.73 days to fruit set, respectively. The minimum days to fruit set (17.3 days) were counted in T₁ (2 g L⁻¹). The higher days to fruit set in T₄ (8 g L⁻¹) might be due to a higher number of pollens, which continued to fertilize for a longer time as compared with lower concentration, which might have pollinated for less time. A lesser number of fruits per bunch was associated with the highest pollen concentration T₄ (8 g L⁻¹), which retained the highest number of fruits per group. The carbohydrates from leaves might have been distributed among the bunches and finally to the fruits. The fewer the fruits, the higher their share of carbohydrates will be and vice versa. Similarly, [Shahid et al. \(2017\)](#) found that when pollination was done at different intervals, the days to fruit set in Dhakki date palms varied.

Table 1: Effect of different suspended pollen concentrations on fruit set percentage and days to fruit set of date palm cultivars Dhakki and Zahidi

Pollen concentration	Fruit set (%)			Days to fruit set		
	Varieties		Mean	Varieties		Mean
	Dhakki	Zahidi		Dhakki	Zahidi	
T ₁ (2 g L ⁻¹)	63.4 7 NS	62.1 3	62.80 d	17.7 9 NS	16.8 3	17.31 d
T ₂ (4 g L ⁻¹)	75.4 7	74.4 9	74.98 c	19.9 9	19.4 6	19.73 c
T ₃ (6 g L ⁻¹)	90.4 7	89.1 8	89.83 b	21.5 9	20.4 5	21.02 b
T ₄ (8 g L ⁻¹)	96.4 7	91.4 6	93.97 a	23.4 0	22.1 0	22.75 a
Mean	81.4 7 NS	79.3 1		20.6 9 NS	19.7 1	

Means having common letters are non-significant at $p < 0.05$

Fruit length (cm)

Data on the result of pollen concentration on the fruit length of date varieties showed significant differences for varieties, pollen concentration and their interaction as well (Table 2). The data in Table 4.3 showed that the maximum fruit length (4.76 cm) was measured in the Dhakki variety, while the minimum 3.87 cm) was noted in Zahidi. The maximum fruit length 5.05 (cm) was recorded in T₁ (2 g/L), followed by T₂ (4 g/L) and T₃ (6 g/L) with 4.55 (cm) and 4.05 (cm), respectively. While the minimum fruit length of 3.61 (cm) was observed in T₄ (8 g/L). The maximum fruit length was noted in Dhakki of 5.63 (cm) with pollen concentration T₁ (2 g/L), followed by Dhakki variety with T₂ (4 g/L) and T₃ (6 g/L) by giving fruit lengths of 5.04 cm and 4.32 cm, respectively. The least fruit weight of 3.87 cm was observed in the Zahidi variety with T₄ (8 g/L), followed by the Zahidi variety with T₃ and T₂, which gave fruit lengths of 3.78 cm and 4.07 cm, respectively. Though the minimum fruit length of 3.17 cm was measured in the Zahidi variety with the highest pollen concentration T₄ (8 g L⁻¹), followed by T₃, T₂ and T₁ (2 g L⁻¹) by producing fruit lengths of 3.78, 4.07 and 4.47 cm, respectively. The maximum fruit length in Dhakki with the lowest pollen concentration (T₁) might be due to a lower number of fruits per group compared with the highest pollen concentration (T₄), which retained the highest number of fruits per bunch. The carbohydrates from leaves might have been distributed among the bunches and finally to the fruits. The fewer the fruits, the higher their share of carbohydrates will be and vice versa. These findings corroborated those of [Abdalla et al. \(2011\)](#) in date palm cv. Zaghloul, who recorded improvement of the fruit physical quality in terms of

increasing the fruit weight and dimensions using pollen grains suspension concentration from 3 to 1.5 g L⁻¹. These findings could be due to the reduction in fruit set because of a lack in the number of fruits per bunch, despite changing the number of leaves, which may induce a better supply of carbohydrates that are manufactured in the leaves.

Fruit width (cm)

Variances for date varieties as pollen concentration, while their interaction was found non-significant (Table 2). Data associated with pollen concentration indicated that maximum fruit width of 2.57 cm was measured in T₁ (2 g L⁻¹), followed by T₂ (4 g L⁻¹) and T₃ (6 g L⁻¹) by producing 2.44 and 2.32 cm fruit width, while the lowest fruit width of 1.96 cm was recommended in T₄ (8 g L⁻¹). Data concerning date varieties showed that Dhakki produced more fruit width (2.56 cm) as associated with Zahidi, which gave a fruit width of 2.09 cm. The maximum fruit width of the Dhakki variety may be due to its genetic characteristics compared to the Zahidi variety. The maximum fruit width in pollen concentration of T₁ (2 g L⁻¹) might be due to a smaller number of fruits which received the maximum share of carbohydrates prepared in the leaves, as compared to T₄ (8 g L⁻¹), which has the highest density of fruit per bunch. These results agreed with the findings of Abdalla et al. (2011) in date palm cv. Zaghloul, who noted improvement of the fruit width in terms of increasing the fruit weight and dimensions by using pollen grain suspension concentration from 3 to 1.5 g L⁻¹. Similarly, Munir et al. (2020) noted maximum fruit width in unlike date palm varieties with spray different pollen concentrations.

Table 2: Effect of different suspended pollen concentrations on fruit length (cm)

and fruit width (cm) of date palm cultivars Dhakki and Zahidi

Pollen concentration	Fruit length (cm)			Fruit width (cm)		
	Varieties		Mean	Varieties		Mean
	Dhakki	Zahidi		Dhakki	Zahidi	
T ₁ (2 g L ⁻¹)	5.63 a	4.47 c	5.0 5 a	2.85 NS	2.28	2.5 7 a
T ₂ (4 g L ⁻¹)	5.04 b	4.07 d	4.5 5 b	2.69	2.19	2.4 4 b
T ₃ (6 g L ⁻¹)	4.32 c	3.78 e	4.0 5 c	2.52	2.13	2.3 2 c
T ₄ (8 g L ⁻¹)	4.04 d	3.17 f	3.6 1 d	2.18	1.74	1.9 6 d
Mean	4.76 a	3.87 b		2.56 NS	2.09	

Means having common letters are non-significant at $p < 0.05$

Fruit weight (g)

Table 4.5 contains information on the weight of the fruits. There were substantial variations in the means for date varieties, pollen concentration, and their interaction, according to the data. The maximum fruit weight, 23.07 (g), was observed in Dhakki, while the minimum, 13.07 (g), was observed in Zahidi. The maximum fruit weight 20.24 (g) was recorded in T₁ (2 g/L), trailed by T₂ (4 g/L) and T₃ (6 g/L) with 19.07 (g) and 17.53 (g), respectively. While the minimum fruit weight of 15.3 (g) was recorded in T₄ (8 g/L). The data about the highest fruit weight of 25.86 (g) was recorded in the Dhakki variety with pollen concentration T₁ (2 g/L), followed by the Dhakki variety with T₂ (4 g/L) and T₃ (6 g/L) by giving pulp weights of 24.24 g and 22.5 g, respectively. The minimum fruit weight of 11.2 g was observed in the Zahidi variety with T₄ (8 g/L), followed by the Zahidi variety with T₃ and T₂, which gave fruit weights of 12.55 g and 13.92 g, respectively. The maximum fruit weight in the case of the Dhakki variety, as compared with the Zahidi variety, might be due to genetic potential, which caused heavy fruits in the Dhakki variety. The heavy fruit

weight of Dhakki with pollen concentration of T_1 (2 g L⁻¹) might be attributed to more available carbohydrates being supplied by leaves to a lesser number of fruits as compared with T_4 (8 g L⁻¹), which retained the maximum number of fruits per plant. Our finding was supported by the previous finding of Abdalla et al. (2011), who reported that the pollen grain suspension at a rate of (1-3 g L⁻¹) increased the fruit weight in date palm cv. Zaghloul.

Pulp weight (g)

Significant variation was observed for pulp weight for date varieties, pollen concentration and their interaction (Table 4.6). The highest pulp weight, 22.06 (g), was recorded in Dhakki, although a minimum of 11.21 (g) was recorded in Zahidi. The maximum pulp weight, 18.75 (g), was recorded in T_1 (2 g/L), followed by T_2 (4 g/L) and T_3 (6 g/L) with 17.81 (g) and 15.40 (g), respectively. While the minimum pulp weight of 14.59 (g) was recorded in T_4 (8 g/L). The data regarding the highest pulp weight of 24.66 (g) was noted in the Dhakki variety with pollen concentration T_1 (2 g/L), followed by the Dhakki variety with T_2 (4 g/L) and T_3 (6 g/L) by giving pulp weights of 23.77 and 20.43 g, respectively. The minimum pulp weight of 9.78 g was measured in the Zahidi variety with T_4 (8 g/L), followed by the Zahidi variety with T_3 and T_2 , which gave pulp weights of 10.38 and 11.84 g, respectively. The higher pulp weight in the case of Dhakki compared with Zahidi might be the result of genetic variation in the size of the fruit. The highest pulp weight of Dhakki with pollen concentration T_1 (2 g L⁻¹) might be due to a reduced number of fruits compared with T_4 (8 g L⁻¹), which has a higher number of fruits for the same available source. Recent studies have demonstrated that pollen grain suspension concentrations significantly influence fruit set percentage, total yield per palm, and various fruit

physical traits, especially pulp weight and fruit mass. For example, Al-Busaidi and Al-Habsi (2006) and Al-Sabahi et al. (2006) found that liquid pollen treatments notably improved reproductive success and physical quality in cultivars like Fard and Khalas. Similarly, El-Salhy (2010) reported that foliar spraying with pollen grain suspensions not only increased fruit set and bunch yield in the Saidy cultivar but also enhanced fruit length, diameter, and pulp-to-seed ratio, confirming the potential of this method in precision pollination strategies.

Table 3: Effect of different suspended pollen concentrations on fruit weight (g) and pulp weight (g) of date palm cultivars Dhakki and Zahidi

Pollen concentration	Fruit weight (g)			Pulp weight (g)		
	Varieties		Mean	Varieties		Mean
	Dhakki	Zahidi		Dhakki	Zahidi	
T_1 (2 g L ⁻¹)	25.8 6 a	14.6 2 e	20.24 a	24.6 6 a	12.8 4 e	18.75 a
T_2 (4 g L ⁻¹)	24.2 4 b	13.9 2 f	19.07 b	23.7 7 b	11.8 4 f	17.81 b
T_3 (6 g L ⁻¹)	22.5 c	12.5 5 g	17.53 c	20.4 3 c	10.3 8 g	15.40 c
T_4 (8 g L ⁻¹)	20.8 3 d	11.2 h	15.3 d	19.3 9 d	9.78 h	14.59 d
Mean	23.0 7 a	13.0 7 b		22.0 6 a	11.2 1 b	

Means having common letters are non-significant at $p < 0.05$ ss

Seed weight (g)

The data in Table 4 showed that significant differences were found for date varieties, pollen concentration and their interaction regarding seed weight. The highest seed weight, 1.29 (g), was recorded in Dhakki, while the minimum, 1.11 (g), was recorded in Zahidi. The maximum seed weight 1.28 (g) was recorded in T_1 (2 g L⁻¹), trailed by T_2 (4 g L⁻¹), T_3 (6 g L⁻¹) with 1.23

and 1.19 (g) respectively. While the minimum seed weight (1.11 g) was recorded in T₄ (8 g L⁻¹). As per the interactive response of pollen concentrations and date palm varieties, the highest seed weight, 1.42 (g), was recorded in T₁ in the Dhakki variety, while the minimum 1.07 (g) seed weight was noted in the Zahidi variety in T₄. The reason for the higher seed weight of Dhakki as compared to the Zahidi variety might be the result of genetic variation. The variation in interaction of the highest seed weight of Dhakki with pollen concentration T₁ (2 g L⁻¹) might be attributed to the bigger size of Dhakki with larger seed weight in case of a small number of fruits per plant, compared with a greater number of fruits in T₄. Our results were supported by Abdalla et al. (2011), who reported that the maximum seed weight (g) was achieved with spraying of (1-3 g L⁻¹) of pollen.

Seed length (cm)

The data shown in Table 4 indicated significant results about seed length of date cultivars for date varieties and pollen concentrations, while their interaction was found non-significant. Data regarding the seed length of date varieties showed that higher seed length (2.79 cm) was observed in the Dhakki variety as compared to the Zahidi variety, which provided a seed length of 2.44 cm. Data about pollen concentration showed that the highest seed length of 2.83 and 2.68 cm was observed in pollen concentration of T₁ (2 g L⁻¹) and T₂ (4 g L⁻¹), while the minimum seed length of 2.51 and 2.47 cm was noted in T₃ (6 g L⁻¹) and T₄ (8 g L⁻¹) pollen concentration treatments. The maximum seed length of the Dhakki variety may be the result of the genetic potential for higher fruit and seed length as compared to the Zahidi variety. The data concerning pollen concentration showed that T₁ and T₂ produced the highest seed length, and it may be attributed to a larger amount of fruit density per bunch as

compared to T₃ (6 g L⁻¹) and T₄ (8 g L⁻¹), which retained the maximum number of fruits per bunch with smaller fruit size. Similar results were quoted by Munir et al. (2020); the maximum seed length was recorded at various levels of pollen spraying.

Table 4: Effect of different suspended pollen concentrations on seed weight (g) and seed length (cm) of date palm cultivars Dhakki and Zahidi

Pollen concent ration	Seed weight (g)			Seed length (cm)		
	Varieties		Me an	Varieties		Me an
	Dha kki	Zah idi		Dha kki	Zah idi	
T ₁ (2 g L ⁻¹)	1.42 a	1.14 d	1.2 8 a	3.00 NS	2.65	2.8 3 a
T ₂ (4 g L ⁻¹)	1.33 b	1.14 d	1.2 3 b	2.89	2.46	2.6 8 a
T ₃ (6 g L ⁻¹)	1.26 c	1.11 de	1.1 9 c	2.65	2.37	2.5 1 b
T ₄ (8 g L ⁻¹)	1.15 d	1.07 e	1.1 1 d	2.56	2.28	2.4 4 b
Mean	1.29 a	1.11 b		2.79 a	2.44 b	

Means having in common. The results are non-significant at $p < 0.05$

Seed width (cm)

Significant response was observed about seed width of date palm cultivars for date palm varieties and pollen concentrations, whereas their interaction was found non-significant (4.9). The higher seed width of 0.86 cm was measured in the Dhakki variety as compared with the Zahidi variety, which produced a seed width of 0.72 cm. The data regarding pollen concentrations indicated that maximum seed width (0.84 cm) was found in T₁ (2 g L⁻¹) followed T₂ (4 g L⁻¹) which gave seed width of 0.80 cm while minimum seed width of 0.77 cm and 0.75 cm was noted in T₃ (6 g L⁻¹) and T₄ (8 g L⁻¹), respectively. The higher seed width of the Dhakki variety as compared to the Zahidi variety might be attributed to the genetic makeup of these varieties. The highest seed width in pollen concentrations T₁ might be due to a smaller

number of fruits with bigger size having thick seeds as compared with T₂ and T₃, which retained a higher number of fruits per bunch with smaller size fruits and seeds. Our outcomes were supported by the previous research of [Munir et al. \(2020\)](#) that the maximum seed width was observed in different levels of pollen spray at different intervals.

Fruit drop percentage

Fruit drop % presented in Table 4.10 depicted significant results regarding date variety, pollen concentration and their interaction. The maximum fruit drop, 38.73, was observed in Dhakki, while the minimum, 22.68, was observed in Zahidi. The maximum fruit drop, 35.27, was noted in T₄ (4 g L⁻¹), trailed by T₃ (6 g L⁻¹) and T₂ (4 g L⁻¹) with 33.27 and 29.53, respectively. While lowest fruit drop, 24.64, was recorded in T₁ (2 g/L). The data about the highest fruit drop of 43.73 was recorded in T₄ (8 g/L) in the Dhakki variety, whereas the minimum 17.41 fruit drop was noted in the variety Zahidi in T₁ (2 g/L). While the lowest fruit drop (17.41%) was recorded in the Zahidi variety with pollen concentration T₂ and T₃ (6 g L⁻¹) by giving (21.74 and 24.78%) respectively. The highest fruit drop % in Dhakki with T₄ might be attributed to heavy fruit holding per plant as compared to pollen concentration T₁ (2 g L⁻¹), in which a smaller number of fruits were maintained due to less fruit set % . The thickly populated fruit per bunch might have pushed the adjacent fruits and caused the dropping of less developed fruits in the bunch. These results are similar to those of [Iqbal et al. \(2011\)](#) and [Shafique et al. \(2011\)](#), who discovered that pollinizers can reduce fruit drop.

Table 5: Effect of different suspended pollen concentrations on seed width (cm) and fruit drop percentage of date palm cultivars Dhakki and Zahidi

Pollen concentration	Seed width (cm)			Fruit drop percentage		
	Varieties		Mean	Varieties		Mean
	Dhakki	Zahidi		Dhakki	Zahidi	
T ₁ (2 g L ⁻¹)	0.93 NS	0.75	0.8 4 a	31.8 6 d	17.4 1 h	24.64 d
T ₂ (4 g L ⁻¹)	0.87	0.73	0.8 0 b	37.3 1 c	21.7 4 g	29.53 c
T ₃ (6 g L ⁻¹)	0.82	0.71	0.7 7 c	41.9 9 b	24.7 8 f	33.39 b
T ₄ (8 g L ⁻¹)	0.80	0.68	0.7 5 c	43.7 4 a	26.8 1 e	35.27 a
Mean	0.86 a	0.72 b		38.7 3 a	22.6 8 b	

Means having common letters are non-significant at $p < 0.05$

Days to maturity

The length of the rising season has a profound effect on the marketability of date palm in different regions. The data regarding days to maturity presented in Table 6 indicated significant variation, whereas date varieties and their interaction were found to be non-significant. The maximum days to maturity (155.10 days) were noted in pollen concentration treatment T₁ (2 g L⁻¹), followed by statistically similar days to maturity in T₂ (4 g L⁻¹), T₃ (6 g L⁻¹) and T₄ (8 g L⁻¹) treatments. The higher days to maturity taken by T₁ (2 g L⁻¹) as compared to other treatments might be due to maximum competition among the fruits for light and nutrients continuous supply for the development of larger and heavier fruits. These findings are consistent with those of [Al-Khalifa \(2006\)](#) and [Iqbal et al. \(2011\)](#), who found that pollen sources influenced fruit ripening time. Similar findings were reported by [Abdel \(2012\)](#).

Fruit weight bunch⁻¹ (kg)

The major yield parameters of date palm are the cluster weight and the number of

bunches per plant. The data presented in Table 6 indicated that date palm varieties and pollen concentrations produced important variation in bunch weight, whereas their interaction could not produce any significant response among the interaction means. The maximum bunch weight of 9.44 kg was noted in the Dhakki variety as compared to the Zahidi variety, which produced a bunch weight of 5.77 kg. The data concerning pollen concentrations indicated that maximum bunch weight of 8.77 kg was noted in T₄ (8 g L⁻¹) followed by statistically like bunch weight of 8.05 and 7.76 kg in T₃ (6 g L⁻¹) and T₂ (4 g L⁻¹), respectively, while the lowest bunch weight of 6.82 kg was observed in T₁ (2 g L⁻¹). The higher single bunch weight of Dhakki as compared to Zahidi variety is a direct response to the genetic potential of the Dhakki variety. The maximum bunch weight in T₄ treatment might be the result of the highest fruit set and retention as compared with the rest of the treatments, which had lower fruit set % and retention. These results agreed with the results of [Awad \(2010\)](#), who observed maximum bunch weight when pollinated with higher concentrations of suspended pollen.

Table 6: Effect of different suspended pollen concentrations on Days to maturity and Fruit weight bunch⁻¹ (kg) of date palm cultivars Dhakki and Zahidi

Pollen concentration	Days to maturity		Fruit weight bunch ⁻¹ (kg)			
	Varieties		Mean	Varieties		Mean
	Dhakki	Zahidi		Dhakki	Zahidi	
T ₁ (2 g L ⁻¹)	155.17 ^{NS}	155.04	155.10 ^a	8.49 ^{NS}	5.16	6.82 ^c
T ₂ (4 g L ⁻¹)	153.73	153.07	153.40 ^b	9.81	5.71	7.76 ^b
T ₃ (6 g L ⁻¹)	153.07	152.67	152.87 ^b	10.32	5.78	8.05 ^b

T ₄ (8 g L ⁻¹)	152.43	152.63	152.53 ^b	11.13	6.41	8.77 ^a
Mean	153.60 ^{NS}	153.35		9.94 ^a	5.77 ^b	

Means having common letters are non-significant at $p < 0.05$

Fruit yield (kg plant⁻¹)

The final goal of all the running performs is to increase the fruit yield of any date palm variety. The significant variation was noted between varieties, pollen concentrations, whereas their interaction was found non-significant (Table 7). The maximum fruit yield plant (159 kg) was observed in Dhakki as compared to the Zahidi variety, which produced a fruit yield of 57.66 kg plant⁻¹. The data with respect to pollen concentrations presented that the highest fruit yield of 121.13 kg per plant was noted in T₄ (8 g L⁻¹), followed by T₃ (6 g L⁻¹) and T₁ (2 g L⁻¹) by producing 111.48 kg and 107.48 kg per plant, respectively. The lowest fruit yield of 93.68 kg per plant was noted in the T₁ (2 g L⁻¹) treatment. The highest fruit yield plant⁻¹ of Dhakki as compared to the Zahidi variety may be due to the genetic potential of high yield as compared to the Zahidi variety. Data about pollen concentrations showed that the highest fruit yield per plant in T₄ (8 g L⁻¹) and T₃ (6 g L⁻¹), as compared to T₂ (4 g L⁻¹) and T₁ (2 g L⁻¹), might be due to heavy fruit set% and retention capacity in T₄ and T₃ treatments. These findings are consistent with those of [Iqbal et al. \(2008\)](#).

Table 7: Effect of different suspended pollen concentrations on the yield of date palm cultivars Dhakki and Zahidi

Pollen concentration	Varieties		Mean
	Dhakki	Zahidi	
T ₁ (2 g L ⁻¹)	135.79 ^{NS}	51.57	93.68 ^c
T ₂ (4 g L ⁻¹)	156.96	57.10	107.48 ^b
T ₃ (6 g L ⁻¹)	165.12	57.83	111.48 ^{ab}
T ₄ (8 g L ⁻¹)	178.13	64.13	121.13 ^a
Mean	159.00 ^a	57.66 ^b	

Conclusion

It was concluded from the present research study that Pollen concentrations (treatments) had varying influences on fruit set, fruit set percent, fruit size, bunch weight, and fruit production per plant, among other things. T₄ (8 g L⁻¹) treatment had the maximum fruit set and fruit production per plant. Treatment T₃ (6 g L⁻¹) and T₂ (4 g L⁻¹) also produced reasonable fruit yield with high-value dates. T₁ (2 g L⁻¹) treatment produced the highest quality fruits, but with a lesser total yield. The effect of pollen concentration was equal and independent on both varieties under study. The highest fruit yield in the Dhakki variety, as compared to the Zahidi variety, may be the result of improved genetic potential of the Dhakki variety.

Conflict of interest

The authors acknowledged that the present study was completed in the absence of any conflict of interest.

References

- Abdalla, M. G., El-Salhy, A. M., & Mostafa, R. A. A. (2011). Effect of some pollination treatments on the fruiting of the Zaghloul date palm cultivar under Assiut climatic conditions. *Assiut J. of Agric. Sci*, 42, 350-362.
- Abdelrahim, M. A., Nooh, M. M., Hassan, W. A., & Rizk, S. M. (2015). Therapeutic potential of date palm pollen for testicular dysfunction induced by thyroid disorders in male rats. *PLoS ONE*, 10(10), e0139493. <https://doi.org/10.1371/journal.pone.0139493>
- Al-Wusaibai, N. A., Ben Abdallah, A., Al-Husainai, M. S., Al-Salman, H., & Elballaj, M. (2012). A comparative study between mechanical and manual pollination in two premier Saudi Arabian date palm cultivars. *Indian Journal of Science and Technology*, 5(4), 2487-2490. <https://doi.org/10.17485/ijst/2012/v5i4.4>
- Al-Busaidi, K., & Al-Habsi, S. E. (2006). Effect of pollen grain suspension on fruit set and physicochemical characteristics of two date palm cultivars. Proceedings of the Conference on Date Palm Production and Processing Technology, Muscat, Oman, May 9-11, p. 18.
- Al-Sabahi, S., Al-Busaidi, K., Al-Amr, S. E., & Al-Abri, S. (2006). Effect of pollen grain suspension on fruit set and yield of date palm (*Phoenix dactylifera*) cultivar Fard. Proceedings of the Conference on Date Palm Production and Processing Technology, Muscat, Oman, May 9-11, p. 18.
- El-Salhy, A. M., El-Bana, A. A., Abdel-Galil, H. A., & Ahmed, E. F. S. (2010, March). Effect of Pollen Grains Suspensions Spraying on Yield and Fruit Quality of 'Saidy' Date Palm. In *IV International Date Palm Conference 882* (pp. 329-336).
- Anonymous. (2005). Seal Management Group. Tissue culture date palm. Retrieved on October 23, 2007 from http://www.sealmanagementgroup.co.uk/z_Date_Palms/tissue_culture_date_palms.htm
- Anonymous. (2006). Weight-loss-advisor.com. Dates are the best nutrition for dieting. Retrieved on October 23, 2007 from http://www.weight-loss-advisor.com/weight_loss_articles/date_nutrition.htm
- Anonymous. (2007). Facts About Fruit.com. Dried fruits. Retrieved on October 23, 2007, from http://www.factsaboutfruit.com/Dried_Fruits.html
- Attalla, A. M., Warring, M. O., & Sharaan, F. A. (1998). Suitable time for two Saudi date palm cultivars. *Alexandria. J. Agric. Res*, 43(3), 203-208.
- Awad, M. A., & Al-Qurashi, A. D. (2012). Partial fruit set failure phenomenon in 'Nabbut-Ali' 'Sabkha' date palm cultivars under a hot arid climate as affected by pollinator type and pollination method. *Scientia horticultrae*, 135, 157-163.
- Haller, J. F., Krawczyk, S. A., Gostilovitch, L., Corkey, B. E., & Zoeller, R. A. (2011). Glucose-6-phosphate isomerase deficiency results in mTOR activation, failed translocation of lipin 1α to the nucleus and hypersensitivity to glucose: Implications for the inherited glycolytic disease. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, 1812(11), 1393-1402.
- Henderson, A. (2009). *Palms of southern Asia* (Vol. 48). Princeton University Press.

- Ibrahim, A. M. F., El-Kobbia, A. M., Kitat, F. M., & Abd El-Kawy, M. M. (1998). Cytogenetic studies on date palm. II. Influence of the storage periods on pollen viability of four Date (*Phoenix dactylifera* L.) male types. *Alexandria Journal of Agricultural Research*, 43(2), 247-254.
- Iqbal, M. A., Sadaqat, H. A., & Khan, I. A. (2008). Estimation of genetic diversity among sunflower genotypes through random amplified polymorphic DNA analysis. *Genet Mol Res*, 7(Suppl 4), 1408-13.
- Iqbal, M., Ghafoor, A., & Rahman, S. (2004). Effect of pollination times on fruit characteristics and yield of date palm cv. Dhakki. *Int. J. Agric. Biol*, 6(1), 96-99.
- Iqbal, M., Ghafoor, A., & Rehman, S. (2004). Evaluation of whorl-wise floral characters of seedling male palms used in pollination of Dhakki in Dera Ismail Khan. *Intern. J. Agric and Bio*, 6(1), 100-107.
- Iqbal, M., Khan, M. N., & Munir, M. (2012). Fruit setting, monthly record of fruit drop, fruit weight and economic yield of Dhakki among six varieties of Date palm. *Sarhad Journal of Agriculture*, 28(1), 23-26.
- Iqbal, M., Khera, R. A., Hussain, T., Sadia, H., Abbas, M., Nazir, A., ... & Younas, U. (2017). Cytotoxicity and bioactivity evaluation of pygmy date palm extracts. *Cur. Sci. Perspect*, 3(2), 106.
- Iqbal, M., Munir, M., & Niamatullah, M. (2011). Physio-chemical characteristics of date palm (*Phoenix dactylifera* L.) cultivars at various maturity stages under environmental conditions of Dera Ismail Khan. *J. Agric. Res*, 49(2), 249-260.
- Jamro, M. M. U. R., Sirohi, M. H., Mirbahar, A. A., Lal, S., & Saand, M. A. (2020). 7. Production and quality of Date fruit in response to pollination methods. *Pure and Applied Biology (PAB)*, 9(3), 1737-1742.
- Maryam, M., Fatima, B., Haider, M. S., Abbas, S., Naqvi, M., Ahmad, R., & Khan, I. A. (2015). Evaluation of pollen viability in date palm cultivars under different storage temperatures. *Pak. J. Bot*, 47(1), 377-381.
- Moshfegh, F., Baharara, J., Namvar, F., Zafar-Balanezhad, S., Amini, E., & Jafarzadeh, L. (2016). Effects of date palm pollen on fertility and development of the reproductive system in female BALB/c mice. *Journal of Herbal Medicine and Pharmacology*, 5(1), 23-28.
- Nasr, T. A., Shaheen, M. A., & Bacha, M. A. (1986). Evaluation of seedling male palms used in pollination in the central region of Saudi Arabia. *Date Palm J*, 4(2), 163-173.
- Omar, A. E. D. K., Ahmed, M. A., & Al-Obeed, R. S. (2015). Improving fruit set, yield and fruit quality of date palm (*Phoenix dactylifera*, L. cv. MNIFI) through bunch spray with boron and zinc. *Journal of Testing and Evaluation*, 43(4), 717-722.
- Samouni, M. T., El-Salhy, A. M., Badawy, I. F., & Ahmed, E. F. (2016). Effect of pollination and thinning methods on yield and fruit quality of Sady Date palms. *Assiut J. Agric. Sci*, 47(3), 92-103.
- Shafique, M., Khan, A. S., Malik, A. U., Shahid, M., Rajwana, I. A., Saleem, B. A., ... & Ahmad, I. (2011). Influence of pollen source and pollination frequency on fruit drop, yield and quality of date palm (*Phoenix dactylifera* L.) cv. Dhakki. *Pakistan Journal of Botany*, 43(2), 831-839.
- Shahid, M. A., Iqbal, M., & Niamatullah, M. (2017). Response of male pollinizers in fruit set, yield and quality of Date palm (*Phoenix dactylifera* L.). Cv. Dhakki. *Sarhad Journal of Agriculture*, 33(1).
- Soliman, S. S., Alebidi, A. I., Al-Saif, A. M., Al-Obeed, R. S., & Al-Bahelly, A. N. (2017). Impact of pollination by pollen grain water suspension spray on yield and fruit quality of Segae date palm cultivar (*Phoenix dactylifera* L.). *Pak. J. Bot*, 49 (1), 119-123.