



Comparative Study of Morphological and Floral Traits in (Rosa Hybrida) cv. "Christian Dior" Cuttings Cultivated on Diverse Growing Media

Usman Shoukat Qureshi¹, Muhammad Noman Asif², Dr Imran Hassan³, Dr Abdul Ahad Qureshi⁴, Muhammad Zeeshan⁵, Muhammad Ziad Abdullah Kayani⁶, Husnain Shamshad⁷, Shakaib Mahmood⁸

¹ Lecturer/Horticulturist, Landscape Studio Lab, Horticulture, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi, Pakistan, Email: nasrkhanpbg@gu.edu.pk (Corresponding Author)

² M.Sc. (Hons) Scholar, Horticulture, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi, Pakistan

³ Chairman/ Associate Professor, Horticulture, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi, Pakistan

⁴ Assistant Professor, Horticulture, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi, Pakistan

^{5,6,7,8} Research fellow, Landscape Studio Lab, Horticulture, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi, Pakistan

Abstract

The "Christian Dior" rose cultivar is known for its large elegant, velvety red-colored scented flowers and excellent shelf life. It has been extensively cultivated globally for its aesthetical, medicinal, and commercial uses. However, media selection also plays a key role in the vegetative and reproductive growth of cut-flower production. Under such conditions, research was conducted to examine the performance of diverse media combinations (T1: S+FYM, T2: C.C+FYM, T3: SN+FYM, T4: S+C.C+FYM, T5: SN+C.C+FYM) on phenotypic and floral attributes of "Christian Dior" rose in thermocol boxes. Results showed that treatment T3 (SN+FYM) shows the maximum plant height of 62.885 cm, number of leaves (121), leaf area (17.342cm²), shoot and root length (8.432cm) (35.765cm), while the least plant height of 25.394 cm was recorded in T1 (S+FYM). Whereas, the maximum shelf life of 12 days, flower and bud diameter (6.416cm)(0.792cm), was recorded in T4 and the least shelf life of 8.5 days was observed in T1 (S+FYM). The treatment T5 (SN+C.C+FYM) takes fewer days, 198 to produce its first flower than T1 (S+FYM) of 238 days. Similarly, results showed that T3 performed significantly better, followed by T5 which produces higher quality cut flowers by providing optimum aeration, essential nutrients, and water-holding ability to enhance its growth and development.

Keywords: Growth substrate, shoot development, rose cultivation, phenotypic traits, organic media, cut flower

DOI: <https://zenodo.org/records/10888892>

Journal Link: <https://jai.bwo-researches.com/index.php/iwr/index>

Paper Link: <https://jai.bwo-researches.com/index.php/iwr/article/view/84>

Publication Process Received: 16 Dec 2024/ Revised: 8 Jan 2025/ Accepted: 27 Jan 2025/ Published: 28 Jan 2025

ISSN: Online [3007-0929], Print [3007-0910]

Copyright: © 2024 by the first author. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Indexing:



Publisher:

BWO Research International (15162394 Canada Inc.) <https://www.bwo-researches.com>

Introduction

Rose (*Rosa hybrida*) cv. “Christian Dior” is a high-quality cutflower cultivar of hybrid tea rose developed in 1958 by Meilland International Nursery, France. This cultivar is a cross between tea roses (*Rosa* × *odorata*) and hybrid perpetual roses (*Rosa* × *damascena*), inheriting all the finest characteristics from both parental species (Ketsa et al., 1993; Taha & Awal, 1994; Singh et al., 2016). Due to its striking red velvety blossoms, elegant fragrance, and long-lasting petals, it holds a substantial reputation in the floral business and is ranked at the top of the floriculture industry (Mirjalili, 2015; Hussen & Yassin, 2013; Jadhav & Gurav, 2018). “Christian Dior” cut roses botanically consist of shrubby deciduous, 2-5 feet tall and 3 to 5 feet wide, dark green glossy foliage with serrated broad leaves, double blooms with 30 to 40 petals in each flower, and stems decked with sharp prickled thorns (Mortazavi et al., 2007; Bhawana et al., 2013). It flourishes best in temperate regions and is spread across Asia, Australia, Europe, and North America. It is also found in limited parts of Africa and South America, where suitable environments favor rose gardening. It grows best on well-drained, loamy to sandy loam soils with neutral to slightly acidic pH 6 to 6.5, requiring a minimum of 6 to 8 hours of direct sunlight daily and temperatures ranging between 15°C and 25°C daily (Sokolov & Petko, 2021). It requires moderate annual rainfall of 500 to 700 mm as high humidity will lead to fungal disease in this cultivar (Ketsa & Dadaung, 2005; Kadam et al., 2023). The success of vegetative growth of *Rosa hybrida* cv. “Christian Dior” cuttings is significantly influenced by the category of media used for growth and rooting. For healthy plant growth and development, an optimal growing medium is required to enhance

propagation efficiency. However, the inadequate comparative study availability on the effect of diverse growing media—such as soil, sand, coconut coir, and their blends creates an opportunity for the researchers (Kaushal & Kumari, 2020). Rose cultivation in alternate growth media except soil is used by the application of sand through nutrient flow method (Takano, 1988; Riffault et al., 2014); gravel culture (Sarro et al., 1989); organic manures mineral wool and aeroponics (Zieslin & Snir, 1989); rockwool (Kool & van de Pol, 1991) and perlite (Katsoulas & Baille, 1999). Aside from soil, the continuing interest in the farming industry is to use the organic by-products as an alternate source of growth that is comparatively cheaper and easily available with low nutritional releases (Mikkelsen, 2003; Ahmad, 2012). Over the last few years, coco coir has been used excessively as an alternate medium due to its many similarities with peat and other factors like acceptable pH, EC, and other chemical properties (Khandaker et al., 2020). Coconut coir has other additional features to be used as a growth medium, including high cation exchange and water-holding capacities, a structure that allows good aeration, and resistance to decomposition (Ahmad et al., 2012). High availability of water, nutrients, and aeration necessary for root development are the important parameters needed to characterize a good soil substrate (Waseem et al., 2013; Altun, 2024). Growth media are important in the growth of plants since they act as a storage of minerals and also facilitate the exchange of gases in plants (Ayesha et al., 2011). The use of organic manure increases the fertility of soils and nutrient supply and improves the growth of plants more than the use of inorganic fertilizers does (Okwuagwu et al., 2003). Potting media for roses must be light,

porous, and well-drained to allow constant harvesting and metabolism of the plants (Khilari, 2021). Peat is extensively used but expensive, which creates a demand for rich alternatives like coconut coir (Dede *et al.*, 2012; Ribeiro *et al.*, 2007). The use of soil-less substrate such as perlite and coconut coir enhances flower yield and quality of stems (Fascella & Zizzo, 2004). Proper rooting and growth of the cuttings can be accomplished by the use of well-aerated and nutrient media (Keisling & Kester, 1979; Monika & Chandla, 2021). This study aimed to assess the impact of different growing media on the rooting capacity and vegetative growth parameters of (*Rosa hybrida*) cv. "Christian Dior" cuttings. The outcomes will aid in the selection of the most appropriate medium that will improve the propagation methods made commercially and horticulturally, thus improving the success rates of practicing rose cultivation at a more economical cost.

Materials and Methods

The present study was conducted at the research field area, Department of Horticulture, Pir Mehr Ali Shah Arid Agriculture University Rawalpindi, Pakistan (2023-2024). A total number of 225 (P×P and R×R= 6 inches) cuttings (8 inches in length) of the rose cultivar "Christian Dior" were collected from Changa Nursery Farm, Mandi Pattoki Punjab, Pakistan. The experiment was laid on thermapol boxes (Dimension: length = 30", width = 18", depth = 10") from the National Institute of Health Islamabad (NIH), and cutting was placed at a 6-inch depth. Then holes were made with the tip of a knife on the bottom of each box for the drainage and aeration of given media. Before plantation, the basic nutrient analysis of media was done at the Soil and Water Testing Laboratory, Data Gunj Bukhsh Road, Rawalpindi, Pakistan (Table 2). The total number of treatments and calculations of different media used is

shown in Table 1 of treatments with three replications.

Procedures for phenotypic and floral attributes

These plants were grown until the full flowering stage. Further phenotypic and floral (plant height, number of leaves, leaf area, fresh weight, dry weight, shoot length, root length, days to flower emergence, flower bud diameter, flower diameter, color variation of flower, shelf life) attributes of plants were recorded according to the standard procedures.

Plant height (cm)

Plant height was measured from the base of the soil to the tip of the plant for 15 plants per treatment in centimeters using a steel measuring scale (model: "NEXT") then, the average was taken.

$$PH^{av} = \frac{(PH1 + PH2 + PH3 + \dots n)}{PHt}$$

PH_{av} stands for average plant height, PH₁ stands for single plant height and PH_t is the total number of plants (Heady, 1957).

Number of leaves (count)

The number of leaves was counted for 5 plants per treatment and their average was calculated using the given formula.

$$NL^{av} = \frac{(NL1 + NL2 + NL3 + \dots n)}{NLt}$$

NL_{av} is the average number of leaves, NL₁ leaves are counted for plants per treatment, and NL_t is the total number of plants.

Leaf area (cm²)

It was measured with the help of a leaf area meter (model: Li-3100) in (cm²) of 20 randomly selected leaves of 15 plants per treatment and their average was calculated.

$$LA^{av} = \frac{(LA1 + LA2 + LA3 + \dots n)}{Lt}$$

LA_{av} stands for average leaf area, LA₁ leaf area of each plant selected, and Lt is the total number of selected plants.

Fresh and dry weight of flower(g)

A weighing balance (model: FA2204) was used to measure both the fresh and dried weights of the flowers in grams (g) for 15 plants per treatment separately and their average was taken.

$$FW/DW^{av} = \frac{\Sigma(FW1+\dots n)\Sigma(DW1+\dots n)}{Nft}$$

Where FW/DW^{av} stands for average fresh and dry weight, Σ (FW1), Σ (DW1) shows the sum of the fresh weight and dry weight of each flower, and N_{Ft} is the total number of flowers (Fanourakis *et al.*, 2020).

Shoot and root length (cm)

The length of the shoot and root was measured in centimeters using a steel measuring scale (model: "NEXT"). The following formula was used to calculate the average length of shoots and roots.

$$SL/RL^{av} = \frac{\Sigma(SL1+\dots n)\Sigma(RL1+\dots n)}{Nt}$$

Where SL/RL^{av} shows the average shoot and root length, Σ (SL1), Σ (SL1) denotes the sum of individual lengths of shoots or roots, and N_t is the total number of plants.

Days to flower emergence (days)

The number of days was calculated from the planting of cuttings till the day of first flower appeared among fifteen flowers from each treatment then the average was taken.

$$DE^{av} = \frac{\Sigma(DE1 + DW2+\dots n)}{Dt}$$

Where DE^{av} stands for the average days to flower emergence, DE1 stands for the day the first flower in a plant.

Flower bud and flower diameter (cm)

Flower and flower bud diameter was measured with the help of an vernier caliper (model: AOS: 500-196-30, Mitutoyo-Japan) in cm of 15 plants per treatment and average was calculated as;

$$FD/FB^{av} = \frac{\Sigma(FD1 + FD2\dots n)\Sigma(FB1 + FB2\dots n)}{Nt}$$

FD/FB^{av} stands for average flower and bud diameter, Σ (FD and FB) refers to the

sum of individual diameters measured, and N_T shows the total number of flowers or flower buds measured (Ma *et al.*, 2005; Nazari *et al.*, 2009).

Color variation of flower (chroma values)

Chromameter was used to understand the color variation in flower that occurs in 5 different treatments. To evaluate color change among various treatments, we calculated the color difference using the formula:

$$\Delta E = \sqrt{(L1 - L2)^2 + (a1 - a2)^2 + (b1 - b2)^2}$$

Where L1, a1, and b1 show the color values from one treatment, L2, a2, and b2 color values from another treatment, and ΔE stands for the total color difference. The higher value of ΔE shows greater color variation.

Shelf life (days)

The shelf life of flowers was recorded by observing the number of days each flower remained healthy and in good condition till its 50% deterioration in distilled water.

$$SL = \frac{(DFD - DIP)}{Nt}$$

Where SL stands for shelf life, DFD is the day when deterioration is first observed, and DIP is the day it when placed in the vase.

Statistical analysis

The data obtained from the experiment was analyzed by using a Randomized Complete Block Design (RCBD) through Analysis of Variance (ANOVA) with the help of Statistix 8.1 software (Steel 1997).

Treatment	Media Combinations	(Ratio = 1:1:1 v/v) in cubic feet per replication box
T ₁	S + FYM	(1.25 + 1.25)
T ₂	C.C + FYM	(1.25 + 1.25)
T ₃	SN + FYM	(1.25 + 1.25)
T ₄	S + C.C + FYM	(0.8333 + 0.8333 + 0.8333)
T ₅	SN + C.C + FYM	(0.8333 + 0.8333 + 0.8333)

Table 1: Different types of growing media with their combinations. (S: Soil, SN=

Sand, FYM= Farm Yard Manure, C.C = Coconut Coir)

Treatments	E.C(dSm ⁻¹)	pH	O.M (%)	P (mg kg ⁻¹)	K (mg kg ⁻¹)	Sat (%)
T ₁ : S + FYM	1.09	7	1.63	5.6	120	55
T ₂ : C.C + FYM	1.21	6.71	4.8	100
T ₃ : SN + FYM	0.97	6.97	1.73	5.8	100	60
T ₄ : S + C.C + FYM	1.37	7.04	1.70	6.2	120	70
T ₅ : SN + C.C + FYM	0.89	7.19	1.79	5.9	140	70

Table 2: Physical and chemical properties of different growing media (before plantation) (S: Soil, SN= Sand, FYM= Farm Yard Manure, C.C = Coconut Coir, E.C = Electrical Conductivity, O.M = Organic Matter, P = Phosphorus, K = Potassium, Sat = Saturation)

Soil and Water Testing Laboratory for Research, Data Ganj Bakhsh Road Rawalpindi, Pakistan.

Results

Plant height

The rose cultivar “*Christian Dior*” showed that treatment T₃ (sand + farm yard manure) attained a maximum plant height of 62.885 cm followed by T₅ (sand + coconut coir + farm yard manure) 46.190 cm. On the other hand, T₂ (coconut coir + farm yard manure) had a plant height of 42.211 cm followed by T₄ (soil + coconut coir + farm yard manure) of 36.749 cm while the least height was obtained in the case of T₁ (soil + farm yard manure) 25.394 cm. The results shown by (T₃) were highly statistically significant ($p < 0.05$) as compared to other treatments as described in Fig.1(a).

Number of leaves

The maximum number of leaves found in T₃ (sand + farm yard manure) is 121.28 followed by plants placed in treatment T₅ (sand + coconut coir + farm yard manure) 93.104. The cuttings placed in media T₄ (soil

+ coconut coir + farm yard manure) were 82.544. Similarly, the number of leaves for rose cutting in T₂ (coconut coir + farm yard manure) was 74.717. The minimum number of leaves was observed in T₁ (soil + farm yard manure) 62.544. From the recorded data it can be concluded that there was a statistically significant ($p < 0.05$) effect of different treatments applied while the results of T₃ were highly significant as shown in Fig 1(b).

Leaf area

The maximum leaf area of 17.342 cm² was observed in T₃ (sand + farm yard manure) followed by T₅ (sand + coconut coir + farm yard manure) 14.954 cm², T₄ (soil + coconut coir + farm yard manure) 13.961 cm², and T₂ (coconut coir + farm yard manure) 13.927 cm² respectively. The lowest possible leaf area was attained by T₁ (soil + farm yard manure) (11.71 cm²). The different treatments showed different results that were quite significant statistically at ($p < 0.05$) as explained in Fig.1(c).

Fresh weight of flower

The maximum fresh weight attained by T₄ (soil + coconut coir + farm yard manure) was 9.2366 g followed by T₂ (coconut coir + farm yard manure) 8.5843 g, T₃ (sand + farm yard manure) 8.5268 g and T₅ (sand + coconut coir + farm yard manure) 8.3852 g. The minimum fresh weight was obtained in the case of T₁ (soil + farm yard manure) 7.0685 g. The results indicated that treatment T₄ was significant while other treatments were non-significant for increasing the fresh weight as given in Fig. 1(d).

Dry weight of flower

The maximum dry weight measured in T₄ (soil + coconut coir + farm yard manure) has 1.4252 g followed by T₂ (coconut coir + farm yard manure) 0.9508 g, T₅ (sand + coconut coir + farm yard manure) 0.8476 g and T₃ (sand + farm yard manure) 0.8404 g.

The minimum dry weight was observed in T₁ (soil + farm yard manure) 0.7564 g. Among all treatments, (T₄) performed better and its results were significant at ($p < 0.05$) as shown in Fig. 1(e).

Shoot Length

The maximum shoot length recorded in T₃ (sand + farm yard manure) was 8.276 cm followed by T₂ (coconut coir + farm yard manure) 7.2293 cm, T₄ (soil + coconut coir + farm yard manure) 6.98 cm, and T₅ (sand + coconut coir + farm yard manure) 6.7573 cm. The minimum shoot length was observed in T₁ (soil + farm yard manure) 6.6427 cm. The treatments applied showed statistically significant results at ($p < 0.05$). Similarly, the result shown by treatment (T₃) was substantial as indicated in Fig. 1(f).

Root length (cm)

The maximum root length observed in T₃ (sand + farm yard manure) was 35.756 cm followed by T₂ (coconut coir + farm yard manure) 29.308 cm, T₄ (soil + coconut coir + farm yard manure) 26.904 cm, and T₅ (sand + coconut coir + farm yard manure) 25.452 cm. In comparison, T₁ (soil + farm yard manure) had a minimum root length that was 19.740 cm. The different treatments when applied significantly affected the root length while (T₃) had significantly better performance as shown in Fig. 1(g).

Days to flower emergence


The treatment T₅ (sand + coconut coir + farm yard manure) has taken a minimum duration of 198.6 days followed by T₄ (soil + coconut coir + farm yard manure) 205.8 days, T₃ (sand + farm yard manure) 211.6 days, and T₂ (coconut coir + farm yard manure) 222.2 days. The maximum time to initiate flowering was taken by T₁ (soil + farm yard manure) of 238.4 days. The treatments have significantly produced different days to initiate flowering while (T₅) significantly minimum days to initiate flowering as described in Fig. 2(a).

Flower bud diameter

The maximum flower bud diameter recorded in T₄ (soil + coconut coir + farm yard manure) was 0.792 cm followed by T₂ (coconut coir + farm yard manure) 0.724 cm, T₅ (sand + coconut coir + farm yard manure) 0.7 cm, and T₃ (sand + farm yard manure) 0.664 cm. The minimum flower bud diameter was measured in T₁ (soil + farm yard manure) 0.62 cm. There is not much difference among the treatments applied except the treatment T₄ (soil + coconut coir + farm yard manure) and T₁ (soil + farm yard manure) a clear difference had been indicated. The results of the treatment (T₄) were significant ($p < 0.05$) among the treatments applied for flower bud diameter as shown in Fig 2(b).

Flower diameter (cm)

The maximum flower diameter observed in T₄ (soil + coconut coir + farm yard manure) was 6.416 cm, followed by T₂ (coconut coir + farm yard manure) 5.432 cm, T₃ (sand + farm yard manure) 5.404 cm, and T₅ (sand + coconut coir + farm yard manure) 5.316 cm. The minimum flower diameter was attained in treatment T₁ (soil + farm yard manure) 5.064 cm. Among all the treatments applied T₄) significantly performed better as shown in Fig. 2(c)











Fig 1. Effect of different growing media on (a) plant height (b) number of leaves (c) leaf area (d) fresh weight of flower (e) dry weight of flower (f) shoot length and (g) root length of rose cutting where different treatments applied shown statistically significant results at (p<0.05). (S: Soil, SN= Sand, FYM= Farm Yard Manure, C.C = Coconut Coir)

Color variation of flower

The chroma meter was used to analyze color variation in flowers across different treatments, with measurements based on light intensity. The maximum lightness value (ΔE) was recorded for T₄ (soil + coconut coir + farm yard manure) with a value of 42.126, followed by T₂ (coconut coir + farm yard manure) with 37.948, T₃ (sand + farm yard manure) with 36.687, and T₅ (sand + coconut coir + farm yard manure) with 29.236. The lowest lightness value was seen in T₁ (soil + farm yard manure) with 24.917. Overall, T₄ showed the most intense color variations with the highest values for both red and yellow hues, while T₁ displayed the least variation in all aspects as shown in Fig. 2(d).

Shelf life

The treatments T₂ (coconut coir + farm yard manure), T₃ (sand + farm yard manure), T₄ (soil + coconut coir + farm yard manure), and T₅ (sand + coconut coir + farm yard manure) show better and same shelf life of cut flower i.e. 9.7 days as compared to T₁ (soil + farm yard manure) which counts 8.5 days of shelf life as shown in Fig. 2(e).


Discussions

The significant increase in plant height noted in T₃ (sand + farmyard manure) can be attributed to the good drainage, aeration, and nutrient supply provided by sand mixed with farmyard manure (Ahmad *et al.*, 2012). Sand-based media increase root development by helping oxygen availability, which supports nutrient uptake and overall plant vigor (Okwuagwu *et al.*, 2003). The lowest performance of T₁ (soil + farmyard manure) may be due to poor compaction and aeration, which inhibits nutrient absorption and root growth (Awang *et al.*, 2009).

The maximum number of leaves recorded in T₃ reflects the suitable physical properties of sand, which help robust root development and nutrient absorption. Sand-based substrates are known to prevent waterlogging, ensuring efficient nutrient uptake and, root respiration (Fascella & Zizzo, 2004). In contrast, T₁'s lesser leaf count may result from reduced compaction in soil and oxygen availability, limiting nutrient uptake and overall vegetative growth (Keisling & Kester, 1979).

The larger leaf area in T₃ aligns with studies that highlight the importance of well-aerated and nutrient-rich media for foliage development (Ahmad *et al.*, 2012). Sand improves root growth by improving air porosity, while farmyard manure supplies essential nutrients, supporting larger leaf development. Conversely, the

minimal leaf area observed in T₁ can be attributed to poor soil aeration, limiting nutrient availability, and leaf expansion (Waseem *et al.*, 2013).






Fig 2. Effect of different growing media on (a) days to flower emergence (b) flower bud diameter (c) flower diameter (d) color variation of flower (e) shelf life of rose cutting where different treatments applied shown statistically significant results at ($p < 0.05$). (S: Soil, SN= Sand, FYM= Farm Yard Manure, C.C = Coconut Coir)

The highest fresh weight of flowers in T₄ (soil + coconut coir + farmyard manure) suggests that coconut coir improves water retention and nutrient availability. Coconut coir is known for its ability to retain nutrients and high water-holding capacity, supporting flower growth (Ribeiro *et al.*, 2007). T₁'s lower fresh weight due to soil compaction, restricting water and nutrient uptake (Dede *et al.*, 2012).

The superior dry weight in T₄ can be attributed to the balanced moisture preservation and aeration properties of coconut coir combined with the nutrient supply from farmyard manure (Fascella & Zizzo, 2004). This combination helps prolonged nutrient availability and optimal plant metabolism. The reduced dry weight in T₁ indicates limited nutrient absorption and poor aeration (Keisling & Kester, 1979).

The maximum shoot length in T₃ is consistent with findings that sandy media improve root aeration and nutrient uptake, helping vegetative growth (Ahmad *et al.*, 2012). Conversely, the minimum shoot length in T₁ reflects the adverse effects of nutrient

absorption and poor soil structure on root development (Awang *et al.*, 2009).

The height root length in T₃ supports the idea that sand increases root elongation by improving drainage and aeration (Fascella & Zizzo, 2004). Suitable oxygen availability in sandy media promotes nutrient absorption and root growth. In contrast, the short root length in T₁ is likely due to poor soil aeration and compaction, hindering root development (Keisling & Kester, 1979).

T₅ showed the minimum time to flower emergence due to the ideal balance of aeration, water retention, and nutrient availability provided by sand, coconut coir, and farmyard manure (Ahmad *et al.*, 2012). In comparison, the late flowering in T₁ due to poor soil structure hinders root development and nutrient uptake, delaying flowering (Tariq *et al.*, 2012).

The big flower bud diameter in T₄ reflects the favorable water retention and nutrient availability provided by coconut coir and farmyard manure (Ribeiro *et al.*, 2007). The bad performance in T₁ is likely due to poor aeration and nutrient uptake (Okwuagwu *et al.*, 2003).

The larger flower diameter in T₄ highlights the importance of balanced moisture retention and nutrient supply for ideal flower development (Awang *et al.*, 2009). Soil compaction in T₁ restricts nutrient absorption, resulting in smaller flowers (Keisling & Kester, 1979).

The highest lightness value in T₄ indicates ideal nutrient availability and water retention, enhancing flower pigmentation (Waseem *et al.*, 2013). Poor soil structure in T₁ results in lesser pigmentation due to nutrient deficiencies (Ahmad *et al.*, 2012).

Treatments T₂, T₃, T₄, and T₅ showed enhanced shelf life due to the balanced nutrient availability and moisture retention provided by coconut coir and sand (Dede *et al.*, 2012). T₁ shorter shelf life is likely due to

poor nutrient supply and moisture preservation (Ribeiro *et al.*, 2007).

Conclusion

In conclusion, on phenotypic characters the T₃ (sand + farm yard manure) performed best due to their natural properties like better soil aeration, best water with holding capacity, and maximum availability of nutrients like freely available phosphorous and potassium in addition to organic contents while, on the other hand in reproductive parameters T₄ (soil + farm yard manure + coconut coir) showed good results because in that treatment media provided a suitable environment for flowering like balanced pH, maximum availability of nutrients with good water holding capacity and improved the physiochemical environment require for optimum flowering. In case of the number of days taken to produce the first flower, the treatment T₅ (sand + farm yard manure + coco peat) performed better because farm yard manure provided the required amount of nutrients and coconut coir added the suitable growth environment by adjusting media pH with increasing the withholding capacity of water besides it also optimized the uptake of nutrients along with farm yard manure. Moreover, the added sand helped to leach down excessive water with other excessive nutrients that were of no use for plant growth.

References

Ahmad, I., Ahmad, A., Ahmad, S., Amjad, A., Saleem, M., & Akram, A. (2012). Effect of Various Agricultural Substrates on Biometric and Qualitative Characteristics of *Ruscus hypophyllum*. *International Journal of Agriculture & Biology*, 14(1).

Altun, B. (2024). Possibilities of Using Organic Wastes as a Growing Medium in Soilless Culture for Cut Flower Rose. *BioResources*, 19(1).

Awang, Y., Shaharom, A. S., Mohamad, R. B., & Selamat, A. (2009). Chemical and physical characteristics of cocopeat-based media mixtures and their effects on the growth and development of *Celosia cristata*. *American*

journal of agricultural and biological sciences, 4(1), 63-71.

- Ayesha, N. R. (2011). Influence of different growth media on the fruit quality and reproductive growth parameters of strawberry (*Fragaria ananassa*). *Journal of Medicinal Plants Research* 5(26).
- Bhawana, L., Ghosal, M., Gupta, S. K., & Mandal, P. (2013). Assessment of Different Preservative Solutions on Vase Life of Cut Roses. *Journal of Ornamental & Horticultural Plants*, 3(3).
- Dede, O. H., Dede, G., & Ozdemir, S. (2012). The use of hazelnut husk and biosolid in substrate preparation for ornamental plants. *African Journal of Agricultural Research*, 7(43), 5837-5841.
- Fanourakis, D., Bouranis, D., Tsaniklidis, G., Rezaei Nejad, A., Ottosen, C.-O., & Woltering, E. J. (2020). Genotypic and phenotypic differences in fresh weight partitioning of cut rose stems: Implications for water loss. *Acta physiologiae plantarum*, 42, 1-10.
- Fascella, G., & Zizzo, G. (2004). Effect of growing media on yield and quality of soilless cultivated rose. *International Symposium on Soilless Culture and Hydroponics*, 697, 133-138.
- Heady, H. F. (1957). The measurement and value of plant height in the study of herbaceous vegetation. *Ecology*, 38(2), 313-320.
- Hussen, S., & Yassin, H. (2013). Review on the impact of different vase solutions on the postharvest life of rose flower. *International Journal of Agricultural Research and Review*, 1(2), 13-17.
- Jadhav, P. B., & Gurav, N. P. (2018). Extending the storage-life and shelf-life of a rose flowers cv. Dutch post-harvest using chemical compounds and cold storage (Ecofrost). *International Journal of Research and Review*, 5(6), 151-156.
- Kadam, G. B., Desai, P., Raju, D. S., Kumar, P. N., Saha, T., Kumar, J., & Prasad, K. (2023). Evaluation of genetic variability in fragrant rose (*Rosa* × *hybrida*) cultivars under open field cultivations. *Current Horticulture*, 11(2), 33-38.
- Katsoulas, C., Kittas, N., & Baille, A. (1999). Transpiration and canopy resistance of greenhouse soilless roses: measurements and modeling. In *III International Workshop on Models for Plant Growth and Control of the Shoot and Root Environments in Greenhouses*, 507, 61-68.
- Kaushal, S., & Kumari, P. (2020). Growing media in floriculture crops. *Journal of Pharmacognosy and phytochemistry*, 9(2), 1056-1061.
- Keisling, T., & Fuqua, M. (1979). Aluminium and manganese toxicity of rose plants grown in East Texas. 509-510.

- Ketsa, S., & Dadaung, S. (2005). *Effect of sodium dichloroisocyanurate and sucrose on vase life of cut roses*. Paper presented at the IV International Symposium on Rose Research and Cultivation 751.
- Ketsa, S., Thampitakorn, F., & Piluek, C. (1993). Effects of silver nitrate and silver thiosulfate on vase-life of cut roses. *Agriculture and Natural Resources*, 27(1), 91-97.
- Khandaker, M. M., Fatini, A., Abdulrahman, M. D., Abdullahi, U. A., & Badaluddin, N. A. (2020). Growing media influence on the growth and development of *Rosa hybrida*. *Plant archives*, 20(2), 6001-6009.
- Khilari, P. (2021). *Effect of different potting media on growth and flowering of spray chrysanthemum (Dendranthema grandiflora Tzvelev) cv. Pusa Guldasta under Malwa Plateau of MP*. Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya.
- Kool, M., & Pol, P. v. d. (1991). The rose cultivar Madelon on rockwool. The rootstock has a considerable influence on flower yield. 62-64.
- Mikkelsen, R. L. (2003). Using Tobacco By-products as a Nitrogen Source for Container-Grown Houseplants. *Journal of plant nutrition*, 26(8), 1697-1708.
- Mirjalili, S. A. (2015). Assessment of concurrent of the sucrose and silver nitrate on cut flower of rose (*Rosa hybrida* cv. 'Red One'). *J. Bio. Env. Sci*, 6, 122-126.
- Monika, K. S. Y., & Chandla, A. (2021). Effect of different potting mixtures on growth and flowering characters of *Chrysanthemum* cv. Haldighati. *Flora and Fauna*, 27(1), 20-26.
- Mortazavi, N., Naderi, R., Khalighi, A., Babalar, M., & Allizadeh, H. (2007). The effect of cytokinin and calcium on cut flower quality in rose (*Rosa hybrida* L.) cv. Illona.
- Nazari, F., Khosh-Khui, M., & Salehi, H. (2009). Growth and flower quality of four *Rosa hybrida* L. cultivars in response to propagation by stenting or cutting in soilless culture. *Scientia Horticulturae*, 119(3), 302-305.
- Okwuagwu, M., Alleh, M., & Osemwota, I. (2003). The effects of organic and inorganic manure on soil properties and yield of okra in Nigeria. *African Crop Science Conference Proceedings*, (6), 390-393.
- Ribeiro, H., Romero, A., Pereira, H., Borges, P., Cabral, F., & Vasconcelos, E. (2007). Evaluation of a compost obtained from forestry wastes and solid phase of pig slurry as a substrate for seedlings production. *Bioresource technology*, 98(17), 3294-3297.
- Riffault, L., Destandau, E., Pasquier, L., André, P., & Elfakir, C. (2014). Phytochemical analysis of *Rosa hybrida* cv. 'Jardin de Granville' by HPTLC, HPLC-DAD and HPLC-ESI-HRMS: Polyphenolic fingerprints of six plant organs. *Phytochemistry*, 99, 127-134.
- Sarro, M., Sanchez, M., Miyar, C., & Zornoza, P. (1988). *Nutritional requirements of two rose cultivars grown in gravel culture*. Paper presented at the International Symposium on Protected Cultivation of Ornamentals in Mild Winter Climates 246.
- Singh, K. P., Bala, M., Panwar, S., & Kumari, S. (2016). Mutation breeding in roses: A review. *Journal of Ornamental Horticulture*, 19(3and4), 55-74.
- Sokolov, T., & Pet'ko, L. (2021). "New Look" And Christian Dior's Roses.
- Taha, R. M., & Awal, A. (1994). Morphogenesis of *Rosa hybrida* Linn var. Christian Dior in vitro. *Malaysian Journal of Science*, 15(1&2), 45-48.
- Takano, T. (1988). *Effect of conductivity and temperature of nutrient solution on the mineral nutrition of horticultural crops in water culture*. Paper presented at the Symposium on High Technology in Protected Cultivation 230.
- Waseem, K., Hameed, A., Jilani, M. S., Kiran, M., Rasheed, M., Javeria, S., & Jilani, T. (2013). Effect of different growing media on the growth and flowering of stock (*Matthiola incana*) under the agro-climatic condition of Dera Ismail Khan. *Pak. J. Agri. Sci*, 50(3), 523-527.
- Zieslin, N., & Snir, P. (1989). Responses of rose plants cultivar 'Sonia' and *Rosa indica* major to changes in pH and aeration of the root environment in hydroponic culture. *Scientia Horticulturae*, 37(4), 339-349.